



Genome Sequence of the *Siphoviridae* *Staphylococcus aureus* Phage vB_SauS_BaqSau1

 Dayan Lozano-Solano,^a Jhonnatan Reales-González,^b Heath W. Catoe,^c Raul R. Raya,^d  Antonio J. Acosta-Hoyos^a

^aUniversidad Simón Bolívar, Facultad de Ciencias Básicas y Biomédicas, Unidad de Genética y Biología Molecular, Barranquilla, Colombia

^bUniversidad del Atlántico, Departamento de Biología, Barranquilla, Colombia

^cUniversity of Miami Miller School of Medicine, Miami, Florida, USA

^dCentro de Referencia para Lactobacilos (CERELA), San Miguel de Tucumán, Argentina

ABSTRACT Here, we report the genome sequence of a *Siphoviridae* phage named vB_SauS_BaqSau1 (BaqSau1), infecting *Staphylococcus aureus*. Phage BaqSau1 was isolated from a sewage water treatment plant in Sahagún, Córdoba, Colombia. It has a double-stranded DNA (dsDNA) genome of 44,384 bp with 67 predicted genes, including a lysin containing a CHAP (cysteine, histidine-dependent amidohydrolase/peptidase) domain.

Staphylococcus aureus is a common pathogen that causes a wide range of infections in humans and animals. The Centers for Disease Control and Prevention (CDC) established the dissemination of methicillin-resistant *Staphylococcus aureus* (MRSA) as a worldwide health problem due to its resistance to multiple antibiotics (www.cdc.gov/drugresistance/index.html), and the World Health Organization (WHO) recently included MRSA in a high-priority list of bacteria for which new antimicrobial strategies are urgently needed (https://www.who.int/medicines/publications/WHO-PPL-Short_Summary_25Feb-ET_NM_WHO.pdf). Several characteristics of phages, for example, specificity, selectivity, self-limiting replication, and constant evolution, make them a promising alternative to antibiotics for treatment of bacterial infections (1). Here, we describe the lysogenic *Staphylococcus aureus* phage vB_SauS_BaqSau1 (BaqSau1).

S. aureus strain RN4220 (2) was used to isolate bacteriophage BaqSau1 from wastewater collected in Sahagún, Córdoba, Colombia, using a 24-hour enrichment process (3), and then to propagate it on tryptic soy (TS) broth or TS agar supplemented with 0.05 mM MgSO₄ using the double-layer overlay technique (4) (Fig. 1A). Genomic DNA was extracted using the PureLink viral RNA/DNA minikit (ThermoFisher Scientific) according to the protocol described by the Center for Phage Technology at Texas A&M University (<https://cpt.tamu.edu/phage-links/phage-protocols/>). Whole-genome sequencing was done by ACGT, Inc. (USA). The DNA was fragmented by ultrasonication to an average target fragment size of 550 bp and used for constructing a sequencing library using the NEXTflex rapid DNA sequencing kit. The final library was sequenced with the Illumina MiSeq v3 flow cell instrument, which generated 6,381,090 300-bp paired-end raw reads. The adapter and low-quality sequences (Q < 30) were trimmed, and short reads (<50 bp) were filtered out using Trimmomatic software v0.36 using default configurations (5). The trimmed reads were *de novo* assembled using SPAdes v3.11.1 (6), resulting in a contig with 86.4× fold coverage. The final contig was annotated with RAST (7) using the *PhiETA2* genome as the reference (taxonomy identification number 326036). The coding DNA sequences (CDSs) and putative functions were predicted by RAST and further analyzed using HHPred (8). Average nucleotide identity (ANI) analysis was done with CLC Genomics Workbench v20.0 (Qiagen) using default configurations.

Citation Lozano-Solano D, Reales-González J, Catoe HW, Raya RR, Acosta-Hoyos AJ. 2020. Genome sequence of the *Siphoviridae* *Staphylococcus aureus* phage vB_SauS_BaqSau1. *Microbiol Resour Announc* 9:e00147-20. <https://doi.org/10.1128/MRA.00147-20>.

Editor John J. Dennehy, Queens College

Copyright © 2020 Lozano-Solano et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Antonio J. Acosta-Hoyos, aacosta24@unisimonbolivar.edu.co.

Received 14 February 2020

Accepted 4 March 2020

Published 9 April 2020




FIG 1 Plaque appearance (A) and virion morphology (B) of bacteriophage vB_SauS_BaqSau1. The plaques were obtained as described in the text. For electron microscopy, a high-titer lysate was applied on Formvar-coated grids, negatively stained with 2% uranyl acetate, and imaged with a Zeiss EM-109 transmission electron microscope (Carl Zeiss AG); the capsid size is approximately 44 nm.

BaqSau1 is a new member of the *Siphoviridae* family (Fig. 1B); its genome consists of a double-stranded DNA molecule of 44,384 bp with an overall G+C content of 34.05%. Average nucleotide identity (ANI) (9, 10) analysis of BaqSau1 with other *Staphylococcus* phages revealed that it has over 90% nucleotide sequence identity with members of the *Phietaivirus* genus (11). Genome analysis revealed a total of 67 coding DNA sequences (CDSs), of which 34 had a predicted function. The functional CDSs were categorized into 4 clusters of functionally related putative genes as follows: (i) phage structure and packaging proteins, 21 CDSs; (ii) DNA replication and regulation proteins, 6 CDSs; (iii) life cycle proteins, 5 CDSs; and (iv) lysis proteins, 2 CDSs. Potential key virulence factors of *S. aureus* were found in 3 CDSs, as well as a CDS encoding a superantigen pathogenicity island SaPI, which provides resistance to beta-lactamases and the Panton-Valentine leukocidin (PVL) toxin gene (12, 13). Open reading frame (ORF) 29 codes for an integrase with a C-terminal integrative and conjugative element from *Bacillus subtilis* (ICEBs)-like catalytic domain (cd01189). The lytic enzymes identified were a class II holin (ORF 26) and an *N*-acetylmuramoyl-L-alanine amidase endolysin (ORF 27) containing a CHAP (cysteine, histidine-dependent amidohydrolase/peptidase) domain (14) similar to LysK from staphylococcal bacteriophage K, a lytic phage against MRSA (15).

Finally, we conclude that BaqSau1 is a new member of the temperate *Siphoviridae* belonging to the *Phietaivirus* genus, containing an endolysin with possible antibacterial activity.

Data availability. The genome sequence of phage BaqSau1 was deposited in GenBank under the accession number [MK658834](https://ncbi.nlm.nih.gov/nucl/MK658834). The raw sequence reads have been submitted to the NCBI SRA under accession number [PRJNA610274](https://ncbi.nlm.nih.gov/sra/PRJNA610274).

ACKNOWLEDGMENT

This work was supported by an internal grant from Universidad Simón Bolívar.

REFERENCES

- Drulis-Kawa Z, Majkowska-Skrobek G, Maciejewska B. 2015. Bacteriophages and phage-derived proteins—application approaches. *Curr Med Chem* 22:1757–1773. <https://doi.org/10.2174/0929867322666150209152851>.

2. Nair D, Memmi G, Hernandez D, Bard J, Beaume M, Gill S, Francois P, Cheung AL. 2011. Whole-genome sequencing of *Staphylococcus aureus* strain RN4220, a key laboratory strain used in virulence research, identifies mutations that affect not only virulence factors but also the fitness of the strain. *J Bacteriol* 193:2332–2335. <https://doi.org/10.1128/JB.00027-11>.
3. Van Twest R, Kropinski AM. 2009. Bacteriophage enrichment from water and soil. *Methods Mol Biol* 501:15–21. https://doi.org/10.1007/978-1-60327-164-6_2.
4. Kropinski AM, Mazzocco A, Waddell TE, Lingohr E, Johnson RP. 2009. Enumeration of bacteriophages by double agar overlay plaque assay. *Methods Mol Biol* 501:69–76. https://doi.org/10.1007/978-1-60327-164-6_7.
5. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
6. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
7. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
8. Zimmermann L, Stephens A, Nam S-Z, Rau D, Kubler J, Lozajic M, Gabler F, Soding J, Lupas AN, Alva V. 2018. A completely reimplemented MPI bioinformatics toolkit with a new HHpred server at its core. *J Mol Biol* 430:2237–2243. <https://doi.org/10.1016/j.jmb.2017.12.007>.
9. Richter M, Rosselló-Móra R. 2009. Shifting the genomic gold standard for the prokaryotic species definition. *Proc Natl Acad Sci U S A* 106:19126–19131. <https://doi.org/10.1073/pnas.0906412106>.
10. Figueras MJ, Beaz-Hidalgo R, Hossain MJ, Liles MR. 2014. Taxonomic affiliation of new genomes should be verified using average nucleotide identity and multilocus phylogenetic analysis. *Genome Announc* 2:e00927-14. <https://doi.org/10.1128/genomeA.00927-14>.
11. Arahal DR. 2014. Whole-genome analyses: average nucleotide identity, p 103–122. *In* Goodfellow M, Sutcliffe I, Chun J (ed), *New approaches to prokaryotic systematics*. Academic Press, Cambridge, MA.
12. Lozano D, Díaz L, Echeverry M, Pineda S, Máttar S. 2010. PVL positive methicillin-resistant *Staphylococcus aureus* (MRSA) strains isolated from healthy individuals in Montería, Córdoba. *Univ Sci (Bogota)* 15: 159–165. (In Spanish.) <https://doi.org/10.11144/javeriana.SC15-2.ppmr>.
13. Alite C, Humphrey S, Donderis J, Maiques E, Ciges-Tomas JR, Penadés JR, Marina A. 2017. Dissecting the link between the enzymatic activity and the SaPI inducing capacity of the phage 80 α dUTPase. *Sci Rep* 7:11234. <https://doi.org/10.1038/s41598-017-11234-9>.
14. Bateman A, Rawlings ND. 2003. The CHAP domain: a large family of amidases including GSP amidase and peptidoglycan hydrolases. *Trends Biochem Sci* 28:234–237. [https://doi.org/10.1016/S0968-0004\(03\)00061-6](https://doi.org/10.1016/S0968-0004(03)00061-6).
15. Becker SC, Foster-Frey J, Donovan DM. 2008. The phage K lytic enzyme LysK and lysostaphin act synergistically to kill MRSA. *FEMS Microbiol Lett* 287:185–191. <https://doi.org/10.1111/j.1574-6968.2008.01308.x>.



Correction for Lozano-Solano et al., “Genome Sequence of the *Siphoviridae* *Staphylococcus aureus* Phage vB_SauS_BaqSau1”

Dayan Lozano-Solano,^{a,e} Jhonnatan Reales-González,^b Heath W. Catoe,^c Raul R. Raya,^d  Antonio J. Acosta-Hoyos^a

^aUniversidad Simón Bolívar, Facultad de Ciencias Básicas y Biomédicas, Unidad de Genética y Biología Molecular, Barranquilla, Colombia

^bUniversidad del Atlántico, Departamento de Biología, Barranquilla, Colombia

^cUniversity of Miami Miller School of Medicine, Miami, Florida, USA

^dCentro de Referencia para Lactobacilos (CERELA), San Miguel de Tucumán, Argentina

^eUniversidad del Valle, Cali, Colombia

Volume 9, no. 15, e00147-20, 2020, <https://doi.org/10.1128/MRA.00147-20>. Page 1:
The author affiliations should appear as shown in this correction.

Citation Lozano-Solano D, Reales-González J, Catoe HW, Raya RR, Acosta-Hoyos AJ. 2020. Correction for Lozano-Solano et al., “Genome sequence of the *Siphoviridae* *Staphylococcus aureus* phage vB_SauS_BaqSau1.” *Microbiol Resour Announc* 9:e01101-20. <https://doi.org/10.1128/MRA.01101-20>.

Copyright © 2020 Lozano-Solano et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Antonio J. Acosta-Hoyos, aacosta24@unisimonbolivar.edu.co.

Published 3 December 2020